JOURNAL OF CHROMATOGRAPHY A

VOL. 781, NOS. 1 + 2 26 SEPTEMBER 1997

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Awareness in Biological Sciences (CABS), Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal, Research Alert and Science Citation Index)

9TH INTERNATIONAL SYMPOSIUM ON HIGH PERFORMANCE CAPILLARY ELECTROPHORESIS AND RELATED MICRO-SCALE TECHNIQUES, ANAHEIM, CA, 26–30 JANUARY 1997

Foreword
by W.S. Hancock (Palo Alto, CA, USA) 1

TECHNIQUES, THEORY AND GENERAL

A simple means of generating pH gradients in capillary zone electrophoresis
by W.E. Rae, J.E. Wong and C.A. Lucy (Calgary, Canada) 3

Effect of the polar groups of anionic surfactant on migration behavior in micellar electrokinetic chromatography
by S. Takeda, S.-I. Wakida, M. Yamane and K. Higashi (Osaka, Japan) and S. Terabe (Hyogo, Japan) 11

Determination of cationic surfactants by capillary electrophoresis with indirect photometric detection
by K. Heinig, C. Vogt and G. Werner (Leipzig, Germany) 17

Prediction of the migration behavior of analytes in capillary electrophoresis based on three fundamental parameters
by P. Britz-McKibbin and D.D.Y. Chen (Vancouver, Canada) 23

Effect of pH and ions in the sample on stacking in capillary electrophoresis
by M.A. Friedberg, M. Hinsdale and Z.K. Shihabi (Winston-Salem, NC, USA) 35

Comparison of polymer coatings of capillaries for capillary electrophoresis with respect to their applicability to molecular imprinting and electrochromatography
by O. Brüggemann and R. Freitag (Ecublens, Switzerland) and M.J. Whitcombe and E.N. Vulfson (Reading, UK) 43

Characterization of inner surface phenomena in capillary electrophoresis capillaries by electron microscopy, atomic force microscopy and secondary ion mass spectroscopy
by S. Kaupp and H. Wätzig (Würzburg, Germany) 55

Quantitative analysis of sodium dodecyl sulphate by capillary electrophoresis
by M.A. Kelly (Bradford, UK), K.D. Altria (Herts, UK) and B.J. Clark (Bradford, UK) 67

Application of capillary electrophoresis–fluorescence line-narrowing spectroscopy for on-line spectral characterization of closely related analytes
by D. Zamzow, C.-H. Lin, G.J. Small and R. Jankowiak (Ames, IA, USA) 73

Determination of kynurenic acid by capillary electrophoresis with laser-induced fluorescence detection
by D.K. Hansen and S.M. Lunte (Lawrence, KS, USA) 81

Interactions of the human class II major histocompatibility complex protein HLA-DR4 with a peptide ligand demonstrated by affinity capillary electrophoresis
by N.H.H. Heegaard, B.E. Hansen, A. Svejgaard and L.H. Fugger (Copenhagen, Denmark) 91

Capillary electrophoresis with electrospray mass spectrometry detection for low-molecular-mass compounds
by T.E. Wheat and K.A. Lilley (Franklin, MA, USA) and J.F. Banks (Branford, CT, USA) 99

Peak shape distortions during capillary electrophoretic separations of multicomponent samples in two co-ion buffers
by R.L. Williams and B. Childs (College Station, TX, USA), E.V. Dose (Winter Park, FL, USA), G. Guiochon (Knoxville, TN, USA) and Gy. Vigh (College Station, TX, USA) 107

Effects of electroosmotic flow on zone mobilization in capillary isoelectric focusing
by Q. Tang and C.S. Lee (Ames, IA, USA) 113

On-line concentration of neutral analytes for micellar electrokinetic chromatography I. Normal stacking mode
by J.P. Quirino and S. Terabe (Hyogo, Japan) 119

Achievement of enantioselectivity of non-polar primary amines by a non-chiral crown ether
by W.X. Huang, S.D. Fazio and R.V. Vivilecchia (East Hanover, NJ, USA) 129

Polyallylamine-supported pseudo-stationary phases for electrokinetic chromatography. Effect of alkyl chain length of the pseudo-stationary phase and methanol content of aqueous buffer on the separation of hydrophobic compounds
by N. Tanaka, K. Nakagawa, H. Iwasaki, K. Hosoya, K. Kimata and T. Araki (Kyoto, Japan) and D.G. Patterson, Jr. (Atlanta, GA, USA) 139

Enantiomer separation of acidic racemates by capillary electrophoresis using cationic and amphoteric β-cyclodextrins as chiral selectors
by Y. Tanaka and S. Terabe (Hyogo, Japan) 151

Dynamics of capillary electrochromatography. Experimental study on the electrosmotic flow and conductance in open and packed capillaries
by G. Choudhary and Cs. Horváth (New Haven, CT, USA) 161

Capillary electrochromatography: theories on electroosmotic flow in porous media (Review)
by A.S. Rathore and Cs. Horváth (New Haven, CT, USA) 185

Automated microanalysis using magnetic beads with commercial capillary electrophoretic instrumentation
by L.G. Rashkovetsky, Y.V. Lyubarskaya, F. Foret, D.E. Hughes and B.L. Karger (Boston, MA, USA) 197

ORGANIC ACIDS

Analysis of organic acids in industrial samples. Comparison of capillary electrophoresis and ion chromatography
by J. Chen, B.P. Preston and M.J. Zimmerman (Newtown Square, PA, USA) 205

Separation of unsaturated fatty acids and related isomeric hydroperoxides by micellar electrokinetic chromatography
by O. Schmitz and S. Gäb (Wuppertal, Germany) 215

Investigation of inorganic ions and organic acids in atmospheric aerosol by capillary electrophoresis
by Z. Krivácsy, Á. Molnár, E. Tarjányi, A. Gelencsér, G. Kiss and J. Hlavay (Veszprém, Hungary) 223

Short chain fatty acids analysis by capillary electrophoresis and indirect UV detection or laser-induced fluorescence (Short communication)
by V. Zuriguel (Ramonville, France), E. Caussé and J.D. Bounéry (Toulouse, France), G. Nouadje, N. Siméon and M. Nertz (Ramonville, France) and R. Salvayre and F. Couderc (Toulouse, France) 233

Comparison of capillary electrophoresis systems for the reliable identification of some carboxylic acids using the marker technique
by S.K. Wiedmer, H. Sirén and M.-L. Riekkola (Helsinki, Finland) and J.H. Jumppanen (Kantvik, Finland) 239

PROTEINS AND ENZYMES

Quaternary structural analysis of nucleoside diphosphate kinases using capillary electrophoresis
by Y.J. Heo and S.Y. Kim (Seoul, South Korea), E. Kim (Taejon, South Korea) and K.-J. Lee (Seoul, South Korea) 251

Application of capillary electrophoresis to determination of enzyme activity and other properties of phosphatidylinositol-specific phospholipase C
by M.J. Suh (Seoul, South Korea), Y.S. Kim (Chungnam, South Korea) and Y.S. Yoo (Seoul, South Korea) 263

Microchip-based capillary electrophoresis of human serum proteins
by C.L. Colyer, S.D. Mangru and D.J. Harrison (Edmonton, Canada) 271

Protein band spreading in capillary zone electrophoresis. Effects of sample zone length and presence of polymer
by S.P. Radko, G.H. Weiss and A. Chrambach (Bethesda, MD, USA) 277

Conventional capillary electrophoresis in comparison with short-capillary capillary electrophoresis and microfabricated glass chip capillary electrophoresis for the analysis of fluorescein isothiocyanate anti-human immunoglobulin G
by I. Rodriguez, Y. Zhang, H.K. Lee and S.F.Y. Li (Singapore, Singapore) 287

NUCLEIC ACIDS AND THEIR CONSTITUENTS

Rapid sizing of polymorphic microsatellite markers by capillary array electrophoresis
by E.S. Mansfield, M. Vainer and D.W. Harris (Sunnyvale, CA, USA), P. Gasparini (Foggia, Italy), X. Estivill (Barcelona, Spain), S. Surrey (Wilmington, DE, USA) and P. Fortina (Philadelphia, PA, USA) 295

Method for detection of epsilon-secondary structure in the precore region of human hepatitis B virus DNA using a fluorescence-based polymerase chain reaction–single-strand-conformation polymorphism technique with capillary electrophoresis
by K. Katsuragi (Tokushima, Japan), K. Kitagishi, T. Mizuguchi and T. Nagashima (Osaka, Japan), M. Kinoshita (Tokushima, Japan) and H. Kumada (Kanagawa, Japan) 307

Separation of DNA sequencing fragments up to 1000 bases by using poly(ethylene oxide)-filled capillary electrophoresis
by Y. Kim and E.S. Yeung (Ames, IA, USA) 315

Application of mixed mobile phases and a step gradient method in capillary electrochromatography for the separation of isomeric polycyclic aromatic hydrocarbon–deoxyribonucleoside adduct mixtures prepared in vitro
by J. Ding (Boston, MA, USA), J. Szeliga and A. Dipple (Frederick, MD, USA) and P. Vouros (Boston, MA, USA) 327

DNA fragment analysis by capillary polymer sieving electrophoresis using poly(acryloylaminoethoxyethanol)-coated capillaries
by K.W. Talmadge, A.K. Tan and M. Zhu (Hercules, CA, USA) 335

Separations of DNA fragments by capillary electrophoresis in N-substituted polyacrylamides
by M. Chiari, S. Riva, A. Gelain, A. Vitale and E. Turati (Milan, Italy) 347

Characterization of TO-PRO-3 as an intercalator for double-stranded DNA analysis with red diode laser-induced fluorescence detection
by S. Rampal, M.-S. Liu and F.-T. A. Chen (Fullerton, CA, USA) 357

Capillary electrophoresis for the detection of known point mutations by single-nucleotide primer extension and laser-induced fluorescence detection
by C.A. Piggee, J. Muth, E. Carrilho and B.L. Karger (Boston, MA, USA) 367

DRUGS, VITAMINS, PHARMACEUTICAL, FORENSIC AND BIOMEDICAL APPLICATIONS

Investigation of the interactions between drugs and mixed bile salt/lecithin micelles. A characterization by micellar affinity capillary electrophoresis (MACE). Part III
by M.A. Schwarz, K. Raith and H.H. Rüttinger (Saale, Germany), G. Dongowski (Rehbrücke, Germany) and R.H.H. Neubert (Saale, Germany) 377

Separation, comparison and identification of fountain pen inks by capillary electrophoresis with UV–visible and fluorescence detection and by proton-induced X-ray emission
by C. Vogt, J. Vogt and A. Becker (Leipzig, Germany) and E. Rohde (Cincinnati, OH, USA) 391

Characterization of metal complexes of pharmaceutical interest by capillary electrophoresis with element sensitive detection
by H. Wittrisch and S. Conradi (Leipzig, Germany), E. Rohde (Cincinnati, OH, USA) and J. Vogt and C. Vogt (Leipzig, Germany) 407

Micellar electrokinetic capillary chromatography for the separation of phenoxymethylpenicillin and related substances
by Y. Zhu, A. Van Schepdael, E. Roets and J. Hoogmartens (Leuven, Belgium) 417

Determination of aspartame by high-performance capillary electrophoresis (Short communication)
by J.J. Pesek and M.T. Matyska (San Jose, CA, USA) 423

Detection of isoforms and isomers of rattlesnake myotoxins by capillary electrophoresis and matrix-assisted laser desorption time-of-flight mass spectrometry
by D. Nedelkov and A.L. Bieber (Tempe, AZ, USA) 429

Determination of L-ascorbic acid in foods by capillary zone electrophoresis
by O.-K. Choi (Seoul, South Korea) and J.-S. Jo (Suwon, South Korea) 435

Quantitation of insulin by capillary electrophoresis and high-performance liquid chromatography. Method comparison and validation
by A. Kunkel, S. Günter, C. Dette and H. Wätzig (Würzburg, Germany) 445

Development of a capillary electrophoresis method with indirect photometric detection for the determination of anions related to anticaries
by T. Wang and S.F.Y. Li (Singapore, Singapore) 457

Anionic–zwitterionic mixed micelles in the micellar electrokinetic separation of clinically relevant steroids on a fused-silica capillary
by G.A. Valbuena, L.V. Rao, J.R. Petersen, A.O. Okorodudu, M.G. Bissell and A.A. Mohammad (Galveston, TX, USA) 467

FOOD AND ENVIRONMENTAL ANALYSIS, OTHER APPLICATIONS

Separation of carnitine and acylcarnitines by capillary electrophoresis (Short communication)
by S. Kiessig and C. Vogt (Leipzig, Germany) 475

Analysis of beverages by capillary electrophoresis (Short communication)
by J.C. Walker, S.E. Zaugg and E.B. Walker (Ogden, UT, USA) 481

Capillary electrophoretic analysis of salicin in *Salix* spp (Short communication)
by S.E. Zaugg, D. Cefalo and E.B. Walker (Ogden, UT, USA) 487

Analysis of nitrate in biological fluids by capillary electrophoresis (Short communication)
by M.A. Friedberg, M.E. Hinsdale and Z.K. Shihabi (Winston-Salem, NC, USA) 491

Capillary electrophoresis and indirect UV detection as a fast and simple analytical tool for bacterial taxonomy (Short communication)
by M. Arellano, S. El Kaddouri, C. Roques, F. Couderc and Ph. Puig (Toulouse, France) 497

Enantiomeric separation of acidic herbicides by capillary electrophoresis using vancomycin as chiral selector
by C. Desiderio, C.M. Polcaro, P. Padiglioni and S. Fanali (Rome, Italy) 503

Study of the enzymatic transformation of fluorescently labeled oligosaccharides in human epidermoid cells using capillary electrophoresis with laser-induced fluorescence detection
by X.C. Le, Y. Zhang, N.J. Dovichi, C.A. Compston, M.M. Palcic, R.J. Beever and O. Hindsgaul (Edmonton, Canada) 515

Identification of antioxidative compounds in plant beverages by capillary electrophoresis with the marker index technique
by A. Kulomaa, H. Sirén and M.-L. Riekkola (Helsinki, Finland) 523

Capillary electrophoresis of cationic and anionic surfactants with indirect conductivity detection
by P.A. Gallagher and N.D. Danielson (Oxford, OH, USA) 533

Capillary electrophoresis as a routine industrial tool for quantitative analytical testing. Determination of sodium dimethyldithiocarbamate in effluents (Short communication)
by A.J. Nitowski (Middlebury, CT, USA), A.A. Nitowski (Wakefield, MA, USA) and J.A. Lent, D.W. Bairley and D. Van Valkenburg (Middlebury, CT, USA) 541

Exoglycosidase matrix-mediated sequencing of a complex glycan pool by capillary electrophoresis
by A. Guttman (San Diego, CA, USA) and K. Williams Ulfelder (Fullerton, CA, USA) 547

Analysis of tetramine in sea snails by capillary electrophoresis–tandem mass spectrometry
by J.-Y. Zhao, P. Thibault, T. Tazawa and M.A. Quilliam (Halifax, Canada) 555

AUTHOR INDEX VOL. 781 565

The table of contents of *Journal of Chromatography* is included in ESTOC – Elsevier Science Tables of Contents service – which can be accessed on the World Wide Web at the following URL addresses:
http://www.elsevier.nl/locate/estoc or http://www.elsevier.com/locate/estoc